## TOPICAL ANAESTHETIC CONSENT FORM



treatment commences, then carefully removed prior to the treatment.  The effects of the anaesthetic vary from individual to individual. We cannot guarantee that you will not experience any discomfort or pain during the treatment and we will not be held liable if the treatment area does not numb.  Allergic reactions can occur from any topical anaesthetics used during the cours of the treatment.  If you do suffer from an allergic reaction during the treatment, you should notify your technician as soon as possible and contact your doctor immediately. If the reaction is very severe you should go to your nearest Accident and Emergency.  An allergic reaction response may display redness, itching, swelling, a rash, blistering, dryness or any other symptom associated with an allergy.  As a result of the permanent cosmetics treatment, combined with the use of the anaesthetic, you can expect to experience swelling and redness that could last between one to five days. You should always follow your post treatment instructions.  For eyeliner treatments you will be asked to keep your eyes closed throughout to numbing period. If for some reason the topical anaesthetic gets into the eye, you must advise your technician at once. The cream will be removed and your eyes to be immediately flushed with a sterile saline solution. It is then safe for the topical anaesthetic to be reapplied.  NOTE: If you experience stinging in the eyes and do not inform your technician immediately, the anaesthetic may numb the eyeball, and a possible corneal abrasion may occur. This can result in a temporary streaming and light sensitivity the eyes. You may be unable to open your eyes and each time you blink it may be painful and temporary blurry vision may occur. Corneal abrasion, however, is rar if you experience any of these symptoms, inform your technician and visit your Doctor immediately.		<ul> <li>It will be applied to the surface of the skin, not injected.</li> </ul>		
of the treatment.  • If you do suffer from an allergic reaction during the treatment, you should notify your technician as soon as possible and contact your doctor immediately. If the reaction is very severe you should go to your nearest Accident and Emergency.  • An allergic reaction response may display redness, itching, swelling, a rash, blistering, dryness or any other symptom associated with an allergy.  • As a result of the permanent cosmetics treatment, combined with the use of the anaesthetic, you can expect to experience swelling and redness that could last between one to five days. You should always follow your post treatment instructions.  • For eyeliner treatments you will be asked to keep your eyes closed throughout the numbing period. If for some reason the topical anaesthetic gets into the eye, you must advise your technician at once. The cream will be removed and your eyes be immediately flushed with a sterile saline solution. It is then safe for the topical anaesthetic to be reapplied.  • NOTE: If you experience stinging in the eyes and do not inform your technician immediately, the anaesthetic may numb the eyeball, and a possible corneal abrasion may occur. This can result in a temporary streaming and light sensitivity the eyes. You may be unable to open your eyes and each time you blink it may be painful and temporary blurry vision may occur. Corneal abrasion, however, is rar if you experience any of these symptoms, inform your technician and visit your Doctor immediately.  have read and fully understood the above and the risks involved with the use of topical anaesthetics and consent to the application and use of the anaesthetic for my permanent cosmetic treatment.	During the treatment	<ul> <li>The effects of the anaesthetic vary from individual to indiviguarantee that you will not experience any discomfort or p</li> </ul>	treatment.  idual. We cannot  iain during the treatment	
your technician as soon as possible and contact your doctor immediately. If the reaction is very severe you should go to your nearest Accident and Emergency.  • An allergic reaction response may display redness, itching, swelling, a rash, blistering, dryness or any other symptom associated with an allergy.  • As a result of the permanent cosmetics treatment, combined with the use of the anaesthetic, you can expect to experience swelling and redness that could last between one to five days. You should always follow your post treatment instructions.  • For eyeliner treatments you will be asked to keep your eyes closed throughout to numbing period. If for some reason the topical anaesthetic gets into the eye, you must advise your technician at once. The cream will be removed and your eyes to be immediately flushed with a sterile saline solution. It is then safe for the topical anaesthetic to be reapplied.  • NOTE: If you experience stinging in the eyes and do not inform your technician immediately, the anaesthetic may numb the eyeball, and a possible corneal abrasion may occur. This can result in a temporary streaming and light sensitivity the eyes. You may be unable to open your eyes and each time you blink it may be painful and temporary blurry vision may occur. Corneal abrasion, however, is rar If you experience any of these symptoms, inform your technician and visit your Doctor immediately.  have read and fully understood the above and the risks involved with the use of topical anaesthetics and consent to the application and use of the anaesthetic for my permanent cosmetic treatment.	Allergic reactions/risks		used during the course	
After the treatment  As a result of the permanent cosmetics treatment, combined with the use of the anaesthetic, you can expect to experience swelling and redness that could last between one to five days. You should always follow your post treatment instructions.  For eyeliner treatments you will be asked to keep your eyes closed throughout to numbing period. If for some reason the topical anaesthetic gets into the eye, you must advise your technician at once. The cream will be removed and your eyes to be immediately flushed with a sterile saline solution. It is then safe for the topical anaesthetic to be reapplied.  NOTE: If you experience stinging in the eyes and do not inform your technician immediately, the anaesthetic may numb the eyeball, and a possible corneal abrasion may occur. This can result in a temporary streaming and light sensitivity the eyes. You may be unable to open your eyes and each time you blink it may be painful and temporary blurry vision may occur. Corneal abrasion, however, is rar If you experience any of these symptoms, inform your technician and visit your Doctor immediately.  have read and fully understood the above and the risks involved with the use of topical anaesthetics and consent to the application and use of the anaesthetic for my permanent cosmetic treatment.		your technician as soon as possible and contact your doctor immediately. If the		
anaesthetic, you can expect to experience swelling and redness that could last between one to five days. You should always follow your post treatment instructions.  • For eyeliner treatments you will be asked to keep your eyes closed throughout to numbing period. If for some reason the topical anaesthetic gets into the eye, you must advise your technician at once. The cream will be removed and your eyes to be immediately flushed with a sterile saline solution. It is then safe for the topical anaesthetic to be reapplied.  • NOTE: If you experience stinging in the eyes and do not inform your technician immediately, the anaesthetic may numb the eyeball, and a possible corneal abrasion may occur. This can result in a temporary streaming and light sensitivity, the eyes. You may be unable to open your eyes and each time you blink it may be painful and temporary blurry vision may occur. Corneal abrasion, however, is rar if you experience any of these symptoms, inform your technician and visit your Doctor immediately.  have read and fully understood the above and the risks involved with the use of topical anaesthetics and consent to the application and use of the anaesthetic for my permanent cosmetic treatment.				
numbing period. If for some reason the topical anaesthetic gets into the eye, you must advise your technician at once. The cream will be removed and your eyes of the immediately flushed with a sterile saline solution. It is then safe for the topical anaesthetic to be reapplied.  • NOTE: If you experience stinging in the eyes and do not inform your technician immediately, the anaesthetic may numb the eyeball, and a possible corneal abrasion may occur. This can result in a temporary streaming and light sensitivity the eyes. You may be unable to open your eyes and each time you blink it may be painful and temporary blurry vision may occur. Corneal abrasion, however, is rar If you experience any of these symptoms, inform your technician and visit your Doctor immediately.  The read and fully understood the above and the risks involved with the use of topical anaesthetics and consent to the application and use of the anaesthetic for my permanent cosmetic treatment.	After the treatment	anaesthetic, you can expect to experience swelling and red between one to five days. You should always follow your p	ness that could last	
immediately, the anaesthetic may numb the eyeball, and a possible corneal abrasion may occur. This can result in a temporary streaming and light sensitivity the eyes. You may be unable to open your eyes and each time you blink it may be painful and temporary blurry vision may occur. Corneal abrasion, however, is rar If you experience any of these symptoms, inform your technician and visit your Doctor immediately.  have read and fully understood the above and the risks involved with the use of topical anaesthetics and consent to the application and use of the anaesthetic for my permanent cosmetic treatment.		numbing period. If for some reason the topical anaesthetic must advise your technician at once. The cream will be rem be immediately flushed with a sterile saline solution. It is the	gets into the eye, you noved and your eyes will	
application and use of the anaesthetic for my permanent cosmetic treatment.		immediately, the anaesthetic may numb the eyeball, and a possible corneal abrasion may occur. This can result in a temporary streaming and light sensitivity of the eyes. You may be unable to open your eyes and each time you blink it may be painful and temporary blurry vision may occur. Corneal abrasion, however, is rare. If you experience any of these symptoms, inform your technician and visit your		
Client name Signature Date	have read and fully understood application and use of the anaes	the above and the risks involved with the use of topical anaesthet hetic for my permanent cosmetic treatment.	ics and consent to the	
	Client name	Signature	Date	